



Image Processing & Pattern

E1425

Lecture 3



Intensity Transformations for Image Enhancement

INSTRUCTOR

DR / AYMAN SOLIMAN

➤ Contents

- Intensity Histogram
- Fixed Intensity Transformations
 - ✓ Negative Transformation
 - ✓ Log Transformation
 - ✓ Power-law (Gamma) Transformation
 - ✓ Thresholding
- Contrast Stretch
 - ✓ Full-Scale Contrast Stretch
- Histogram Equalization
 - ✓ Cumulative Histogram
 - ✓ Intermediate Image




➤ Intensity transformation

- Also called point operation
- Zero-memory operation

➤ Map a given gray level to another level

$$s = T(r)$$

typically, monotonically increasing (but not always)




Intensity Histogram


➤ Intensity Histogram

- Example
a 4x4, 4bits/pixel image →


1	8	6	6
6	3	11	8
8	8	9	10
9	10	10	7



➤ Intensity Histogram






From [Gonzalez & Woods]



➤ Intensity Histogram

From [Gonzalez & Woods]




Fixed Intensity Transformations

➤ Basic Transformations

FIGURE 3.3 Some basic gray-level transformation functions used for image enhancement.

From [Gonzalez & Woods]



Negative:

$$s = L - 1 - r$$

Log:

$$s = c \log(1 + r)$$

Inverse Log:

$$s = e^{cr} - 1$$


Power-law:

$$s = cr^\gamma$$

.....

➤ Negative Transformation

$$s = L - 1 - r$$



a b

FIGURE 3.4
(a) Original digital mammogram.
(b) Negative image obtained using the negative transformation in Eq. (3.2-1).
(Courtesy of G.E. Medical Systems.)


From [Gonzalez & Woods]

➤ Log Transformation

a b

FIGURE 3.5

(a) Fourier spectrum.
(b) Result of applying the log transformation given in Eq. (3.2-2) with $c = 1$.



From [Gonzalez & Woods]

➤ Power-law (Gamma) Transformation

$$s = cr^\gamma$$

From [Gonzalez & Woods]




FIGURE 3.6 Plots of the equation $s = cr^\gamma$ for various values of γ ($c = 1$ in all cases).

➤ Power-law (Gamma) Transformation

$$s = cr^\gamma$$

From [Gonzalez & Woods]



a
b
c
d


FIGURE 3.8
(a) Magnetic resonance (MR) image of a fractured human spine.
(b)–(d) Results of applying the transformation in Eq. (3.2-3) with $c = 1$ and $\gamma = 0.6, 0.4$, and 0.3 , respectively. (Original image for this example courtesy of Dr. David R. Pickens, Department of Radiology and Radiological Sciences, Vanderbilt University Medical Center.)

➤ Power-law (Gamma) Transformation

$$s = cr^\gamma$$


a b
c d

FIGURE 3.9
(a) Aerial image.
(b)–(d) Results of
applying the
transformation in
Eq. (3.2-3) with
 $c = 1$ and
 $\gamma = 3.0, 4.0,$ and
 $5.0,$ respectively.
(Original image
for this example
courtesy of
NASA.)




From [Gonzalez & Woods]

➤ Thresholding



$$s = \begin{cases} 0 & \text{if } r \leq m \\ c & \text{if } r > m \end{cases}$$

m : threshold



➤ Example: Fixed Intensity Transformation

- A 4x4, 4bits/pixel image

1	8	6	6
6	3	11	8
8	8	9	10
9	10	10	7

an intensity transformation

$$s = T(r) = \text{round} \left(\frac{1}{15} r^2 \right)$$

$$1 \rightarrow \text{round}(0.0667) = 0;$$

$$3 \rightarrow \text{round}(0.6) = 1;$$

$$6 \rightarrow \text{round}(2.4) = 2;$$

$$7 \rightarrow \text{round}(3.2667) = 3;$$

$$8 \rightarrow \text{round}(4.2667) = 4;$$

$$9 \rightarrow \text{round}(5.4) = 5;$$

$$10 \rightarrow \text{round}(6.6667) = 7;$$


$$11 \rightarrow \text{round}(8.0667) = 8;$$

passes through

The resulting image is:



0	4	2	2
2	1	8	4
4	4	5	7
5	7	7	3

➤ Example: Histogram Change



Contrast Stretch

➤ General Idea: Make Best Use of the Dynamic Range



➤ Contrast Stretch

General form:

$$s = \begin{cases} \frac{s_1}{r_1} \cdot r & 0 \leq r < r_1 \\ \frac{s_2 - s_1}{r_2 - r_1} \cdot r + \frac{s_1 r_2 - s_2 r_1}{r_2 - r_1} & r_1 \leq r \leq r_2 \\ \frac{2^B - 1 - s_2}{2^B - 1 - r_2} \cdot r + (2^B - 1) \cdot \frac{s_2 - r_2}{2^B - 1 - r_2} & r_2 < r \leq 2^B - 1 \end{cases}$$

Special case → Full-scale contrast stretch:

$$r_1 = r_{\min}$$

$$s_1 = 0$$



$$s = (2^B - 1) \cdot \frac{r - r_{\min}}{r_{\max} - r_{\min}}$$

$$r_2 = r_{\max}$$

$$s_2 = 2^B - 1$$

Typically used: $s = \text{round} \left((2^B - 1) \cdot \frac{r - r_{\min}}{r_{\max} - r_{\min}} \right)$

➤ Example: Full-Scale Contrast Stretch

- Full-scale contrast stretch of a 4×4 , 4bits/pixel image
- Find when $r_{\min} = 4$ $r_{\max} = 11$ $2^B - 1 = 15$

$$s = \text{round} \left((2^B - 1) \cdot \frac{r - r_{\min}}{r_{\max} - r_{\min}} \right)$$

$4 \rightarrow \text{round}(0) = 0;$

$6 \rightarrow \text{round}(4.29) = 4;$

$7 \rightarrow \text{round}(6.43) = 6;$

$8 \rightarrow \text{round}(8.57) = 9;$

$9 \rightarrow \text{round}(10.71) = 11;$

$10 \rightarrow \text{round}(12.86) = 13;$


$11 \rightarrow \text{round}(15) = 15;$

The resulting image is:

4	8	6	6
6	4	11	8
8	8	9	10
8	11	10	7


0	9	4	4
4	0	15	9
9	9	11	13
9	15	13	6

➤ Example: Histogram Change





Histogram Equalization

➤ Histogram Equalization



From [Gonzalez & Woods]

➤ Histogram Equalization



From [Gonzalez & Woods]

➤ Example

- A 4x4, 4bits/pixel image
- First try full-scale contrast stretch $r_{\min}=2$ $r_{\max}=11$

$$s = \text{round} \left((2^B - 1) \cdot \frac{r - r_{\min}}{r_{\max} - r_{\min}} \right)$$


2 → round(0) = 0;
3 → round(1.67) = 2;
8 → round(10.00) = 10;
9 → round(11.67) = 12;
10 → round(13.33) = 13;
11 → round(15) = 15;

The resulting image is:

2	8	9	9
2	3	10	9
8	3	3	11
8	3	10	11

0	10	12	12
0	2	13	12
10	2	2	15
10	2	13	15


➤ Example: Histogram Change



➤ Cumulative Histogram

2	8	9	9
2	3	10	9
8	3	3	11
8	3	10	11

k	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
H(k)	0	0	2	4	0	0	0	0	3	3	2	2	0	0	0	0
Q(k)	0	0	2	6	6	6	6	6	9	12	14	16	16	16	16	16



➤ Intermediate Image

k	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
$H(k)$	0	0	2	4	0	0	0	0	3	3	2	2	0	0	0	0
$Q(k)$	0	0	2	6	6	6	6	6	9	12	14	16	16	16	16	16
original	2	8	9	9	2	3	10	9	8	3	3	11	8	3	10	11
	2	9	12	12	2	6	14	12	9	6	6	16	9	6	14	16



intermediate image

➤ Full-Scale Contrast Stretch of Intermediate Image

intermediate image

2	9	12	12
2	6	14	12
9	6	6	16
9	6	14	16

$$r_{\min}=2$$

$$r_{\max}=16$$

$$s = \text{round} \left((2^B - 1) \cdot \frac{r - r_{\min}}{r_{\max} - r_{\min}} \right) = \text{round} \left(15 \cdot \frac{r - 2}{16 - 2} \right) = \text{round} \left(\frac{15}{14} (r - 2) \right)$$

$$2 \rightarrow \text{round}(0) = 0;$$

$$6 \rightarrow \text{round}(4.29) = 4;$$

$$9 \rightarrow \text{round}(7.50) = 8;$$

$$12 \rightarrow \text{round}(10.71) = 11;$$

$$14 \rightarrow \text{round}(12.86) = 13;$$

$$16 \rightarrow \text{round}(15) = 15;$$

result:

histogram equalized image

0	8	11	11
0	4	13	11
8	4	4	15
8	4	13	15

➤ Histogram Comparison

4	8	6	6
6	4	11	8
8	8	9	10
8	11	10	7


original

0	10	12	12
0	2	13	12
10	2	2	15
10	2	13	15

direct full-scale contrast stretch


0	8	11	11
0	4	13	11
8	4	4	15
8	4	13	15

histogram-equalized



more equalized

➤ Summary of the Histogram Equalization Algorithm



Thank
you

